arXiv:1102.4798v1 [math.DG] 23 Feb 2011

CONVERGENCE OF KÄHLER-RICCI FLOW ON FANO MANIFOLDS, II

GANG TIAN AND XIAOHUA ZHU*

Abstract. In this paper, we give an alternative proof for the convergence of Kähler-Ricci flow on a Fano mnaifold (M, J). This proof differs from that in [TZ3]. Moreover, we generalize the main theorem of [TZ3] to the case that (M, J) may not admit any Kähler-Einstein metrics.

1. Introduction

In this paper, we use Perelman's W-functional in [Pe] to study the convergence of Kähler-Ricci flow on a Fano mnaifold. We want to prove

Theorem 1.1. *Let (M, J) be a compact Kähler-Einstein manifold with positive first Chern class $c_1(M) > 0$. Then for any initial Kähler metric g with Kähler class $2\pi c_1(M)$, Kähler-Ricci flow converges to a Kähler-Einstein metric in the C^∞-topology. Moreover, the convergence can be made exponentially modulo a holomorphism transformation on (M, J).*

Theorem 1.1 was first announced by G. Perelman when he was visiting MIT in the Spring of 2003. In [TZ3], we gave a proof of Theorem 1.1 by using an inequality of Moser-Trudinger type established in [T1] and [TZ1][1]. The main purpose of this paper is to give an alternative proof of this theorem. This new proof does not use the inequality of Moser-Trudinger type. Moreover, our proof also yields a generalization as follows:

Theorem 1.2. *Let (M, J) be a compact Kähler manifold with positive first Chern class $c_1(M) > 0$. Suppose that there exists a Kähler metric g_0 with Kähler class $2\pi c_1(M)$ such that Kähler-Ricci flow with the initial metric g_0 converges to a Kähler-Einstein metric g_{KE} in the C^∞-topology, which may not be compatible with J. Then for any initial Kähler metric g with Kähler class $2\pi c_1(M)$, the Kähler-Ricci flow converges to g_{KE} in the C^∞-topology.*

1991 *Mathematics Subject Classification.* Primary: 53C25; Secondary: 53C55, 58E11.

Key words and phrases. Kähler-Ricci flow, Kähler-Einstein metric, Perelman's W-functional .

* Partially supported by the NSFC Grant 10990013.

[1] We need to add more details about how to use the Moser-Trudinger typed inequality in the case that M has non-trivial holomorphic fields, .

Theorem 1.2 implies that the convergence of Kähler-Ricci flow is independent of the choice of initial Kähler metrics with Kähler class $2\pi c_1(M)$. The main idea in the proof of Theorem 1.1 and Theorem 1.2 comes from [TZ4], where we used Perelman's W-functional to study the stability of Kähler-Ricci flow on a Fano manifold. Recently Sun and Wang obtained a more general result for the stability [SW]. Our proof depends on certain asymptotic estimates for the minimizing functions (we call f_t-functions) defined by the W-functional associated to evolved metrics g_t of Kähler-Ricci flow. Our estimates depend only on the lower bound of Mabuchi's K-energy on the space of Kähler potentials on (M, J) [Ma]. If (M, J) is a Kähler-Einstein manifold, Bando and Mabuchi proved that the K-energy is bounded from below ([BM], also see [DT]). Recently, in [Ch], Chen gives a generalization of this. In fact, Chen's result will be used in the proof of Theorem 1.2.

The organization of this paper is as follows: In Section 2, we outline an approach of proof to Theorem 1.1. In Section 3, we recall the W-functional and discuss some of its applications to Kähler-Ricci flow (cf. Proposition 3.3 and Lemma 3.4). In Section 4, we give some estimates for f_t-functions along the Kähler-Ricci flow. Theorem 1.1 and Theorem 1.2 will be proved in Section 6 and Section 7, respectively. In Section 5, a stability result for Kähler-Ricci flow, Proposition 2.1, will be proved. Section 8 is an appendix in which we give a C^0-estimate for f_t-functions.

Acknowledgements. Part of the work for this paper was done when both authors were visiting the Simons Center for Geometry and Physics at Stony Brook in early 2011. The authors would like to thank people there for hospitality and excellent working condition.

2. An approach of continuity method to Kähler-Ricci flow

Let (M, J) be an n-dimensional compact Kähler manifold with positive first Chern class $c_1(M) > 0$ and g be a Kähler metric with its Kähler class equal to $2\pi c_1(M)$. We consider the (normalized) Kähler-Ricci flow (abbreviated as KR-flow):

$$\frac{\partial g(t)}{\partial t} = -\mathrm{Ric}(g(t)) + g(t), \ g(0) = g. \tag{2.1}$$

It was proved in [Ca] that (2.1) has a global solution $g_t = g(t)$ for all time $t > 0$. For simplicity, we denote by $(g_t; g)$ a solution of (2.1) with initial metric g.

Now we assume that (M, J) admits a Kähler-Einstein metric ω_{KE} with Kähler class $2\pi c_1(M)$.

Proposition 2.1. *Let (M, J) be a Kähler-Einstein manifold with positive first Chern class $c_1(M) > 0$ and g_{KE} be a Kähler-Einstein metric on (M, J)*

with its Kähler form $\omega_{g_{KE}} \in 2\pi c_1(M)$. *Then there exists a number* $\delta > 0$ *which depends only on* g_{KE} *such that for any Kähler metric* g *on* (M, J) *with Kähler form* $\omega_g \in 2\pi c_1(M)$ *satisfying*

$$\|g - g_{KE}\|_{C^3(M)} \leq \delta, \tag{2.2}$$

KR-flow $(g_t; g)$ *converges to* g_{KE} *globally in the* C^∞*-topology.*

Proposition 2.1 is a special case of more general stability theorem on Kähler-Ricci flow on a Fano manifold, which was recently proved by Sun and Wang by using an inequality of Lojasiewicz type for the Kähler-Ricci flow [SW]. We note that Proposition 2.1 was also proved by the second named author if the condition (2.2) is replaced by certain condition on Kähler potentials [Zhu]. In this paper, for the readers' convenience, we will include a proof of Proposition 2.1 in Section 5. Similar argument will be used in proving Theorem 1.2. Here we denote by $\|g - g'\|_{C^\ell(M)}$ the norm:

$$\|g - g'\|_{C^\ell(M)} = \inf_{\Phi}\{|g - \Phi^*(g')|_{C^\ell(M)}\},$$

where Φ runs over all diffeomorphisms of M and $|g_1 - g_2|_{C^\ell(M)}$ denotes the C^ℓ-norm between two tensors g_1 and g_2 on M. The convergence of Riemannian metrics in the C^∞-topology means the convergence in all C^ℓ-norms $\|\cdot\|_{C^\ell(M)}$.

We can write

$$\omega_g = \omega_\phi = \omega_{g_{KE}} + \sqrt{-1}\partial\overline{\partial}\phi \in 2\pi c_1(M)$$

for a Kähler potential ϕ on (M, J) and we define a path of Kähler forms

$$\omega_s = \omega_{g_{KE}} + s\sqrt{-1}\partial\overline{\partial}\phi.$$

Set

$$I = \{s \in [0,1] \mid (g_t^s; \omega_s) \text{ converges to } g_{KE} \text{ in the } C^\infty - \text{topology}\}.$$

Clearly, it follows from Proposition 2.1 that I is not empty. We want to show that I is both open and closed. It follows that $I = [0, 1]$. Theorem 1.1 will follow from this.

The openness of I follows from the following corollary of Proposition 2.1:

Corollary 2.2. *Let* $(M, J, \omega_{g_{KE}})$ *be a Kähler-Einstein manifold with* $\omega_{g_{KE}} \in 2\pi c_1(M) > 0$. *Suppose that there exists a Kähler metric* g_0 *on* (M, J) *with Kähler class* $2\pi c_1(M)$ *such that KR-flow* $(g_t; g_0)$ *converges to* g_{KE} *in the* C^∞*-topology. Then there exists a* $\delta > 0$ *which depends only on* g_{KE} *and* g_0 *such that for any Kähler metric* g *on* (M, J) *with Kähler class* $2\pi c_1(M)$ *satisfying*

$$\|g - g_0\|_{C^3(M)} \leq \delta, \tag{2.3}$$

KR-flow $(g_t; g)$ *converges to* g_{KE} *in the* C^∞*-topology.*

Proof. By the assumption, we can choose T sufficiently large such that if $\tilde{g}_t$ is the solution of (2.1) with initial metric g_0, then

$$\|\tilde{g}_T - g_{KE}\|_{C^3(M)} < \frac{\delta}{2},$$

where δ is a small number determined in Proposition 2.1. Since the Kähler-Ricci flow is stable for any fixed finite time, there is a small $\epsilon > 0$ such that whenever $\|g - g_0\|_{C^3(M)} < \epsilon$, we have

$$\|g_T - \tilde{g}_T\|_{C^3(M)} < \frac{\delta}{2}.$$

Hence, we have

$$\|g_T - g_{KE}\|_{C^3(M)} < \delta. \tag{2.4}$$

Then the flow $(g_t; g_T)$ with initial g_T will converge to g_{KE} in the C^∞-topology according to Proposition 2.1. This proves the corollary. □

It remains to prove

Theorem 2.3. *I is closed.*

This will be proved in Section 6.

3. PERELMAN'S W-FUNCTIONAL

In this section, we review Perelman's W-functional in [Pe]. The W-functional is defined for triples (g, f, τ) on a given closed manifold M of dimension m, where g is a Riemannian metric, f is a smooth function and τ is a constant. It is defined as follows:

$$W(g, f, \tau) = (4\pi\tau)^{-m} \int_M [\tau(R(g) + |\nabla f|^2) + f - m] e^{-f} dV_g, \tag{3.1}$$

where $R(g)$ denotes the scalar curvature of g. We also normalize the triple (g, f, τ) by

$$(4\pi\tau)^{-m} \int_M e^{-f} dV_g = (4\pi\tau)^{-m} V. \tag{3.2}$$

In our situation, since M is a compact Kähler manifold of complex dimension n, we further have the following normalization of the volume of g:

$$\int_M dV_g = (2\pi)^n \int_M c_1(M)^n \equiv V. \tag{3.3}$$

Note that $m = 2n$. Then the W-functional depends only on a pair (g, f) and can be reexpressed as follows:

$$\begin{aligned} W(g,f) &= W(g,f,\frac{1}{2}) + (2\pi)^{-m} mV \\ &= (2\pi)^{-m} \int_M [\frac{1}{2}(R(g) + |\nabla f|^2) + f] e^{-f} dV_g, \end{aligned} \tag{3.4}$$

where (g, f) satisfies (3.2) with $\tau = \frac{1}{2}$ and (3.3). Note that (3.3) holds automatically for certain geometric problems, for example, metrics which evolve along a normalized Ricci flow.

Following Perelman, we define an entropy $\lambda(g)$ by

$$\lambda(g) = \inf_f \{W(g,f) |\ f \text{ satisfies } (3.2)\}.$$

It is well-known that $\lambda(g)$ can be attained by some f (cf. [Ro]). In fact, such a f satisfies the Euler-Lagrange equation of $W(g,\cdot)$,

$$\triangle f + f + \frac{1}{2}(R - |\nabla f|^2) \ = \ \lambda(g). \tag{3.5}$$

Following a computation in [Pe], we can deduce the first variation[2] of $\lambda(\cdot)$,

$$\delta\lambda(g) \ = \ -\int_M < \delta g, \mathrm{Ric}(g) - g + \nabla^2 f > e^{-f} dV_g, \tag{3.6}$$

where $\mathrm{Ric}(g)$ denotes the Ricci tensor of g and $\nabla^2 f$ is the Hessian of f. It follows from (3.6) that g is a critical point of $\lambda(\cdot)$ if and only if g is a gradient shrinking Ricci soliton, namely, g satisfies

$$\mathrm{Ric}(g) \ + \ \nabla^2 f \ = \ g, \tag{3.7}$$

where f is a minimizer of $W(g,\cdot)$.

Next we will prove the uniqueness of solutions for (3.5) if g is a gradient shrinking Ricci soliton.

Lemma 3.1. *If g satisfies* (3.7)*, then any solution of* (3.5) *is equal to the function f in* (3.7) *modulo a constant. Consequently, the minimizer of $W(g,\cdot)$ is unique if the metric g is a gradient shrinking Ricci soliton.*

Proof. Let σ_t be an one-parameter subgroup generated by the gradient field

$$X = \frac{1}{2}\nabla_g f = \frac{1}{2} g^{ij} f_j \frac{\partial}{\partial x_i}.$$

---

[2] It is not obvious that $\lambda(\cdot)$ is differentiable. This can be proved by using the comparison principle and the regularity results we have on (3.5) (see Lemma 3.5 below and Appendix in Section 8). Here we just compute its first variation at those g where λ is smooth. This is the case when the minimizer f is unique. It is sufficient for us in this paper.

Put $\tilde{g}(t,\cdot) = \sigma_t^\star g$. Then $\tilde{g}(t,\cdot)$ is a family of Ricci solitons and $\tilde{g}$ satisfies the normalized Ricci flow,

$$\frac{\partial \tilde{g}}{\partial t} = L_{(\sigma_t^{-1})_\star X}\tilde{g} = -\mathrm{Ric}(\tilde{g}) + \tilde{g}.$$

Let f' be another solution of (3.5) and ρ_t be an one-parameter subgroup generated by the gradient field

$$X' = \frac{1}{2}\nabla_{\tilde{g}} f'.$$

Set $g' = \rho_t^\star \tilde{g}(t,\cdot)$. Then g' satisfies a modified Ricci flow:

$$\begin{aligned}\frac{\partial g'}{\partial t} &= -\mathrm{Ric}(g') + g' - L_{(\rho_t^{-1})_\star X'} g' \\ &= -\mathrm{Ric}(g') + g' - \nabla^2 \rho_t^\star f'.\end{aligned}$$

Clearly, $(\sigma_t \circ \rho_t)^\star f'$ satisfies (3.5) with $\hat{g}_t := (\sigma_t \circ \rho_t)^\star g$ in place of g. Therefore, using the first variation of the functional W and its invariance under diffeomorphisms, we have

$$\begin{aligned}0 &= \frac{dW((\sigma_t \circ \rho_t)^\star g, (\sigma_t \circ \rho_t)^\star f')}{dt} \\ &= -\int_M < \frac{d\hat{g}_t}{dt}, \mathrm{Ric}(\hat{g}_t) - \hat{g}_t + \nabla^2(\sigma_t \circ \rho_t)^\star f' > e^{-(\sigma_t\circ\rho_t)^\star f'} dV_{\hat{g}_t} \\ &= \int_M |\mathrm{Ric}(\hat{g}_t) - \hat{g}_t + \nabla^2(\sigma_t \circ \rho_t)^\star f'|^2 \, e^{-(\sigma_t\circ\rho_t)^\star f'} dV_{\hat{g}_t}.\end{aligned}$$

It follows

$$\mathrm{Ric}(g) - g + \nabla^2 f' = 0.$$

Hence, $f' = f + \mathrm{const.}$. □

As a converse to Lemma 3.1, we have

Lemma 3.2. *Let g be a gradient shrinking Ricci soliton and f be the function in* (3.7) *for g. Then f satisfies* (3.5)*.*

Proof. Differentiating (3.7) and applying the second Bianchi identity, we have

$$\nabla_i R = 2R_{ij} f_j.$$

It follows

$$\nabla_i R + 2f_{ij} f_j - 2f_i = 2(R_{ij} + f_{ij} - g_{ij}) f_j = 0.$$

This implies

$$R + |\nabla f|^2 - 2f \equiv const,$$

and consequently, by using (3.7), we deduce that f is a solution of (3.5). □

Proposition 3.3. *Let* (M, J, g_{KE}) *be a Kähler-Einstein manifold with Kähler class* $2\pi c_1(M)$. *Then* g_{KE} *is a global maximizer of* $\lambda(\cdot)$ *in the space of Kähler metrics* (g, J') *on* M *with Kähler class* $2\pi c_1(M)$.

Proof. By Lemma 3.1, the minimizer f of $W(g_{KE}, \cdot)$ must be a constant. As a consequence, $f = 0$ because of the normalization condition (3.2). Thus for any Kähler metric g with Kähler class $2\pi c_1(M)$, we have

$$\begin{aligned}\lambda(g_{KE}) \,&=\, W(g_{KE}, 0) \,=\, W(g, 0)\\ &\geq\, \inf\{W(g, f) \mid \int_M e^{-f} dV_g \,=\, V\} \,=\, \lambda(g).\end{aligned}$$

Here we have used the fact:

$$\begin{aligned}W(g, 0) \,&=\, n(2\pi)^{-2n} \int_M \mathrm{Ric}(g) \wedge \omega_g^{n-1}\\ &=\, n(2\pi)^{-2n} \int_M \mathrm{Ric}(g_{KE}) \wedge \omega_{g_{KE}}^{n-1} \,=\, n(2\pi)^{-2n} V.\end{aligned}$$

□

The following is a direct corollary of Proposition 3.3 by using the Kähler-Ricci flow.

Lemma 3.4. *Let* (J_0, g_{KE}) *and* (J, g) *be a Kähler-Einstein metric and a Kähler metric on* M *with* $2\pi c_1(M)$ *as both Kähler classes. Suppose that* $\lambda(g) \geq \lambda(g_{KE})$. *Then* g *is a Kähler-Einstein metric with respect to the complex structure* J.

Proof. Let $g(t, \cdot)$ be a solution of (2.1) with the initial metric g. Then by the monotonicity of $\lambda(g)$ along the Ricci flow [Pe] (see (3.6) or (3.8) below), we have

$$\lambda(g(t, \cdot)) \geq \lambda(g), \ \forall\ t > 0.$$

Thus by Proposition 3.3, we see

$$\lambda(g(t, \cdot)) = \lambda(g_{KE}), \ \forall\ t > 0.$$

Again by (3.6), it follows that $g(t, \cdot)$ are all gradient shrinking Ricci solitons. Moreover, the Kähler condition implies that $g(t, \cdot)$ are in fact all Kähler-Ricci solitons. Therefore, by the uniqueness of Kähler-Ricci solitons [TZ2], we see that $g(t, \cdot)$ are all the same modulo automorphisms of M, and consequently, g is a Kähler-Ricci soliton with respect to some holomorphic field X induced by a Hamiltonian function f which satisfies (3.5). On the other hand,

$$\lambda(g) = W(g, f) = W(g, 0) = \lambda(g_{KE}).$$

This implies that $f = 0$ since the minimizer of $W(g, \cdot)$ is unique according to Lemma 3.1. Hence, g is a Kähler-Einstein metric with respect to the complex structure J.

□

We will end this section by a well-known fact, which follows from the monotonicity of Perelman's entropy $\lambda(\cdot)$. For the readers' convenience, we include its proof.

Lemma 3.5. *Let $g_t = g(t,\cdot)$ be a solution of* (2.1) *on M. Suppose that there exists a sequence g_i of g_t converging to a limit Riemannian metric g_∞ in the C^3-topology. Then g_∞ is a Kähler-Ricci soliton with respect to some complex structure J_∞.*

Proof. By the assumption, there exists a sequence of diffeomorphisms Ψ_i of M such that

$$|\Psi_i^\star g_i - g_\infty|_{C^3(M)} \to 0.$$

It follows that after taking a subsequence if necessary, $(\Psi_i^{-1})_\star \cdot J \cdot (\Psi_i)_\star$ converges to an integral complex structure J_∞ in the C^3-topology. The standard regularity theory implies that J_∞ is in fact analytic. By (3.6), we have

$$\frac{d}{dt}\lambda(g_t) = \int_M |\sigma_t^\star(\mathrm{Ric}(g_t) - g_t + \nabla^2 f_t)|^2_{\sigma_t^\star g} dV_{\sigma_t^\star g}, \tag{3.8}$$

where f_t are minimizing solutions of (3.5) associated to metrics g_t and σ_t is the family of diffeomorphisms of M generated by the time-dependent vector field $\frac{1}{2}\nabla_{g_t} f_t$. Since

$$\lambda(g_t) \le W(g_t, 0) = (2\pi)^{-m}\frac{m}{2}V$$

is bounded from above, there exists a sequence of α_i such that as i tends to ∞, $|\alpha_i - i| \to 0$ and

$$\int_M |\mathrm{Ric}(g_{\alpha_i}) - g_{\alpha_i} + \nabla^2 f_{\alpha_i}|^2_{g_{\alpha_i}} dV_{g_{\alpha_i}} \to 0. \tag{3.9}$$

Since the Ricci flow depends continuously on initial metrics, we may choose $\delta_0 > 0$ such that

$$|\Psi_i^\star g_{\alpha_i} - g_\infty|_{C^3(M)} \le \delta_0 \tag{3.10}$$

as $i \to \infty$. On the other hand, by Proposition 8.1 in our Appendix, we know that f_t are uniformly bounded. Thus by applying the regularity theory of elliptic equations to (8.1) in Appendix and using (3.10), we get

$$\|\Psi_i^\star f_{\alpha_i}\|_{C^{4,\frac{1}{2}}(M)} \le C$$

for some uniform constant C. Hence, by taking a subsequence if necessary, there exists a $C^{4,\frac{1}{2}}$-smooth function f_∞ such that

$$\Psi_{\alpha_i}^\star f_{\alpha_i} \to f_\infty$$

in the C^4-topology. Then it follows from (3.9)

$$\mathrm{Ric}(g_\infty) - g_\infty + \nabla^2 f_\infty \;=\; 0.$$

By the regularity theory, f_∞ is actually smooth. Furthermore, since g_∞ is Kähler, the gradient vector field ∇f_∞ has to be holomorphic. This proves that g_∞ is a Kähler-Ricci soliton.

$\square$

4. Estimates for f-function

In this section, we derive some a priori estimates for the f_t-functions solving (3.5) associated to the KR-flow (2.1). Since the flow preserves the Kähler class, we may write the Kähler form of g_t as

$$\omega_\phi \;=\; \omega_g + \sqrt{-1}\partial\overline{\partial}\phi$$

for some Käher potential $\phi = \phi_t$. Furthermore, as usual, one can choose ϕ_t appropriately such that (2.1) is reduced to a parabolic complex Monge-Ampère flow for $\phi(t,\cdot) = \phi_t$,

$$\frac{\partial\phi}{\partial t} = \log\frac{\det(g_{i\bar{j}} + \phi_{i\bar{j}})}{\det(g_{i\bar{j}})} + \phi - h',\;\; \phi(0,\cdot) = 0, \tag{4.1}$$

where $h' = -\frac{\partial\phi}{\partial t}|_{t=0}$ is a Ricci potential of the initial metric g. It is easy to check that each $\frac{\partial\phi}{\partial t}$ is a Ricci potential of g_t. We set $h_t = -\frac{\partial\phi}{\partial t} + c_t$ for some constant c_t so that

$$\int_M e^{h_t}\omega_{g_t}^n \;=\; \int_M \omega_g^n. \tag{4.2}$$

The following estimates are due to G. Perelman. We refer the readers to [ST] for their proof.

Lemma 4.1. *There are constants c and C depending only on the initial metric g such that (a) diam$(M, g_t) \le C$; (b) vol$(B_r(p), g_t) \ge cr^{2n}$; (c) $\|h_t\|_{C^0(M)} \le C$; (d) $\|\nabla h_t\|_{g_t} \le C$; (e) $\|\Delta h_t\|_{C^0(M)} \le C$.*

Under the assumption that the K-energy is bounded from below, we can improve the above estimates as follows.

Proposition 4.2. *If in addition the K-energy is bounded from below on the space of Kähler potentials on (M, J) in Lemma 4.1. Then we have:*
(a) $\lim_{t\to\infty}\|h_t\|_{C^0(M)} = 0$*;*
(b) $\lim_{t\to\infty}\|\nabla h_t\|_{g_t} = 0$*;*
(c) $\lim_{t\to\infty}\|\Delta h_t\|_{C^0(M)} = 0$.

Proof. Recall that the K-energy $\mu(\cdot)$ is defined for a Kähler potential ϕ on (M,J) by

$$\mu(\phi)=-\int_0^1\int_M\dot{\psi}(R(\psi_s)-n)\omega_{\psi_s}^n,$$

where ψ_s is a path of Kähler potentials on (M,J) and $R(\psi_s)$ denote scalar curvatures of Kähler forms ω_{ψ_s}. Then for a family of Kähler potentials $\phi=\phi_t$ in (4.1), we have

$$\frac{d\mu(\phi)}{dt}=-\int_M|\nabla h_t|_{g_t}^2\omega_{g_t}^n.$$

Thus by the lower bound of $\mu(\cdot)$, one sees that there exists a sequence of $t_i\in[i,i+1]$ such that

$$\lim_{t_i\to\infty}H(t_i)=\lim_{t_i\to\infty}\int_M|\nabla h_{t_i}|_{g_{t_i}}^2\omega_{g_{t_i}}^n=0. \tag{4.3}$$

Since $H(t)$ satisfies a differential equation,

$$\frac{dH(t)}{dt}\le CH(t),$$

for some uniform constant (cf. [CT2], [PSSW]), it is easy to derive

$$\lim_{t\to\infty}\int_M|\nabla h_t|_{g_t}^2\omega_{g_t}^n=0. \tag{4.4}$$

Let

$$\tilde{h}_t\ =\ h_t-\frac{1}{V}\int_M h_te^{h_t}\omega_{g_t}^n.$$

By using the weighted Poincare inequality in [TZ3], we have

$$\int_M\tilde{h}_t^2e^{h_t}\omega_{g_t}^n\le\int_M|\nabla h_t|_{g_t}^2e^{h_t}\omega_{g_t}^n.$$

It follows from (c) in Lemma 4.1,

$$\lim_{t\to\infty}\int_M\tilde{h}_t^2\omega_{g_t}^n=0. \tag{4.5}$$

We claim

$$\lim_{t\to\infty}\|\tilde{h}_t\|_{C^0(M)}=0. \tag{4.6}$$

So we get (a) in our proposition by the normalization condition (4.2). In fact the claim follows from an inequality

$$\|\tilde{h}_t\|_{C^0(M)}^{n+1}\le C\|\nabla h_t\|_{g_t}^n[\int_M\tilde{h}_t^2\omega_{g_t}^n]^{\frac{1}{2}}. \tag{4.7}$$

(4.7) can be proved by using the non-collapsing estimate (b) in Lemma 4.1 (cf. [PSSW], [Zh]). By (d) in Lemma 4.1, the claim follows.

Next, set

$$a_t = -\int_M \frac{\partial\phi}{\partial t}\omega_{g_t}^n,$$

a direct computation shows

$$\frac{da_t}{dt} = a_t - \int_M |\nabla h_t|_{g_t}^2 \omega_{g_t}^n.$$

It follows that

$$e^{-t}a_t - a_0 = \int_0^t e^{-s}|\nabla h_s|_{g_s}^2\omega_{g_s}^n \wedge ds.$$

On the other hand, since the K-energy is bounded from below, as in [CT1] or [TZ3], we can normalize h' in (4.1) by adding a suitable constant so that

$$\int_M h'\omega_g^n = -\int_0^\infty \int_M |\nabla h_t|_{g_t}^2 e^{-t}\omega_{g_t}^n \wedge dt.$$

Thus we get

$$a_t = \int_t^\infty e^{t-s}|\nabla h_s|_{g_s}^2\omega_{g_s}^n \wedge ds \to 0, \text{ as } t \to \infty. \tag{4.8}$$

Hence, by (a) in our proposition, we also get

$$\lim_{t\to\infty} \|\frac{\partial\phi}{\partial t}\|_{C^0(M)} = 0. \tag{4.9}$$

Now (b) and (c) follow from the standard gradient and Laplacian estimates for the Kähler-Ricci flow which can be derived by using the Maximum principle as in [Ba], [T1]. For instance, the following lemma was proved in [PSSW] and can be applied to complete the proof of (b) and (c).

□

Lemma 4.3. *([PSSW]) There exist $\delta, K > 0$ depending only on the dimension n with the following property: For any $\epsilon > 0$ with $\epsilon < \delta$ and any $t > 0$, if*

$$\|\frac{\partial\phi}{\partial t}\|_{C^0(M)} \le \epsilon,$$

then

$$\|\nabla h_{t+2}\|_{g_{t+2}} + \|\Delta h_{t+2}\|_{C^0(M)} \le K\epsilon.$$

From Proposition 4.2, we can deduce

Proposition 4.4. *Suppose that the K-energy is bounded from below. Then there exists a sequence of $t_i \in [i, i+1]$ such that*
(a) $\lim_{t_i\to\infty} \|\Delta f_{t_i}\|_{L^2(M,\omega_{g_{t_i}})} = 0$*;*
(b) $\lim_{t_i\to\infty} \|\nabla f_{t_i}\|_{L^2(M,\omega_{g_{t_i}})} = 0$*;*
(c) $\lim_{t_i\to\infty} \int_M f_{t_i}e^{-f_{t_i}}\omega_{g_{t_i}}^n = 0.$

Proof. Let σ_t be the family of diffeomorphisms generated by the time-dependent gradient vector field $\frac{1}{2}\nabla_{g_t} f_t$. Then by (3.6), we have

$$\begin{aligned}\frac{d}{dt}\lambda(g_t) &= \int_M \sigma_t^\star(|\mathrm{Ric}(g_t) - g_t + \nabla^2 f_t|^2 e^{-f_t})[\sigma_t^\star(\omega_{g_t})]^n,\\ &= \int_M |\mathrm{Ric}(g_t) - g_t + \nabla^2 f_t|^2_{g_t} e^{-f_t}\omega^n_{g_t}.\end{aligned}$$

It follows from (2.1) that

$$\begin{aligned}\frac{d}{dt}\lambda(g_t) &= \int_M |\frac{1}{2}\partial\overline{\partial}h_t - D^2 f_t|^2_{g_t} e^{-f_t}\omega^n_{g_t}\\ &\geq \frac{1}{2n}\int_M |\Delta(f_t - h_t)|^2 e^{-f_t}\omega^n_{g_t}.\end{aligned}$$

Since $\lambda(g_t) \leq (2\pi)^{-n} nV$ are uniformly bounded, we see that there exists a sequence of $t_i \in [i, i+1]$ such that

$$\lim_i \int_M |\triangle(f_{t_i} - h_{t_i})|^2 e^{-f_{t_i}}\omega^n_{g_{t_i}} = 0.$$

Thus by Proposition 4.2, we will get (a). Here we used the fact that f_t are uniformly bounded according to Proposition 8.1 in Appendix. Moreover, by

$$\int_M |\nabla f_t|^2 \omega^n_{g_t} = -\int_M f_t \triangle f_t \omega^n_{g_t} \leq C\left(\int_M |\Delta f_t|^2 \omega^n_{g_t}\right)^{\frac{1}{2}},$$

Thus we get (b) from (a).

Define

$$\tilde{f}_t = f_t - \frac{1}{V}\int_M f_t e^{h_t}\omega_t^n.$$

By using the weighted Poincare inequality in [TZ3], we have

$$\int_M \tilde{f}_t^2 e^{h_t}\omega^n_{g_t} \leq 2\int_M |\nabla f_t|^2 e^{h_t}\omega^n_{g_t} \leq C[\int_M |\Delta f_t|^2 \omega^n_{g_t}]^{\frac{1}{2}}.$$

Hence, by using (a) and the boundedness of h_t, we can deduce

$$\int_M \tilde{f}^2_{t_i}\omega^n_{g_{t_i}} \to 0, \text{ as } t_i \to \infty. \tag{4.10}$$

Since

$$-\triangle\tilde{f}_t = -\Delta f_t = f + \frac{1}{2}(R - |\nabla f|^2) - \lambda(g) \leq C',$$

we have

$$-\Delta\tilde{f}_t^+ \leq C',$$

where $\tilde{f}_t^+ = \max\{\tilde{f}_t, 0\}$. By using the standard Moser's iteration, we can derive

$$\tilde{f}_t^+ \leq C''\|\tilde{f}_t^+\|_{L^2(M,g_t)} \leq C''\|\tilde{f}_t\|_{L^2(M,g_t)}.$$

Thus by (4.10), we get

$$\lim_{t_i\to\infty}\|\tilde f_{t_i}^{+}\|_{C^0(M)} = 0. \tag{4.11}$$

Let $\tilde f_t^{-} = \min\{\tilde f_t, 0\}$. Since f_t is uniformly bounded and h_t converges to 0, it follows from $\int_M \tilde f_t e^{h_t}\omega_{g_t}^n = 0$

$$\lim_{t_i\to\infty}\int_M \tilde f_{t_i}^{-}\omega_{g_{t_i}}^n = 0.$$

Thus

$$\begin{aligned} |\int_M \tilde f_{t_i} e^{-\tilde f_{t_i}}\omega_{g_{t_i}}^n| &\le \int_{M^+} \tilde f_{t_i}^{+} e^{-\tilde f_{t_i}}\omega_{g_{t_i}}^n + \int_{M^-}(-\tilde f_{t_i}^{-})e^{-\tilde f_{t_i}}\omega_{g_{t_i}}^n\\ &\le \|\tilde f_{t_i}^{+}\|_{C^0(M)}\int_M e^{-\tilde f_{t_i}}\omega_{g_{t_i}}^n + \sup_M e^{-\tilde f_{t_i}}\int_M(-\tilde f_{t_i}^{-})\omega_{g_{t_i}}^n\\ &\to 0,\ \text{as } t_i\to\infty. \end{aligned} \tag{4.12}$$

Here we denote $M^+ = \{\tilde f_{t_i}\ge 0\}$ and $M^- = \{\tilde f_{t_i}\le 0\}$ and we used the fact that $\tilde f$ are uniformly bounded. By (4.12) and the normalization condition (3.2), we will get the property (c) since f_t differs from $\tilde f_t$ by a function $c(t)$ with $\lim_{i\to\infty} c(t_i) = 0$.

□

Definition 4.5. *The energy level $L(g)$ of entropy $\lambda(\cdot)$ along the KR-flow $(g_t; g)$ with an initial Kähler metric g is given by*

$$L(g) \;=\; \lim_{t\to\infty}\lambda(g_t).$$

By the monotonicity of $\lambda(g_t)$, we see that $L(g)$ exists and it is finite. In case that the K-energy is bounded from below, from Proposition 4.4 together with Proposition 4.2, we get

Corollary 4.6. *Suppose that the Mabuchi's K-energy is bounded from below. Then*

$$L(g) \;=\; (2\pi)^{-n} nV \;=\; \sup\{\lambda(g')|\ \omega_{g'}\in 2\pi c_1(M)\}.$$

The above corollary shows that the energy level $L(g)$ of entropy $\lambda(\cdot)$ does not depend on the initial Kähler metric g.

5. Proof of Proposition 2.1

The proof of Proposition 2.1 depends on the following uniqueness result recently proved by Chen and Song in [CS]. This more general uniqueness has been conjectured by the first named author in [T2]. Note that the uniqueness of Kähler-Einstein metrics on a Fano manifold was proved by Bando and Mabuchi in 1985 for a fixed complex structure [BM].

Theorem 5.1. *([CS]) Let (M, J) be a compact Kähler manifold with $c_1(M) > 0$. Suppose that there are two sequences of Kähler metrics with Kähler class $2\pi c_1(M)$, which converge to Kähler-Einstein metrics (g_1, J_1) and (g_2, J_2) in the C^∞-topology, respectively. Then (g_1, J_1) is conjugate to (g_2, J_2) by a diffeomorphism of M.*

Remark 5.2. *In [CS], Chen and Sun also showed a version of Theorem 5.1 for Kähler metrics with constant scalar curvature and integral Kähler class.*

Remark 5.3. *In fact, we only need a much weaker version of Theorem 5.1 in order to prove Proposition 2.1. Such a weaker version can be proved by using the deformation theory.*

Proof of Proposition 2.1. Let $g_t = g(t, \cdot)$ be a family of evolved Kähler metrics of KR-flow $(g_t; g)$. We claim that there exists $\delta > 0$ such that if g satisfies (2.2), then

$$\|g_t - g_{KE}\|_{C^3(M)} \le \epsilon(\delta), \ \forall \ t > 0, \tag{5.1}$$

where $\epsilon(\delta) \to 0$ as $\delta \to 0$. We will use an argument by contradiction from [TZ4]. On contrary, we may find a sequence of Kähler metrics g^i on (M, J) with Kähler forms in $2\pi c_1(M)$ and a sequence of diffeomorphisms Ψ_i on M which satisfy

$$|\Psi_i^\star g^i - g_{KE}|_{C^3(M)} \to 0, \tag{5.2}$$

such that there exists a number $\epsilon_0 > 0$ and a sequence of Kähler metrics $g^i(t_i, \cdot)$ with property:

$$\|g^i_{t_i} - g_{KE}\|_{C^3(M)} \ge \epsilon_0, \tag{5.3}$$

where $g^i_{t_i}$ is a solution of KR-flow $(g^i_t; g^i)$ at time t_i. Moreover, by the stability of Ricci flow in finite time and by the continuity of the Ricci flow, one can choose the number ϵ_0 arbitrary small in (5.1) and can further assume that $g^i_{t_i}$ satisfies

$$2\epsilon_0 \ge \|g^i_{t_i} - g_{KE}\|_{C^3(M)} \ge \epsilon_0. \tag{5.4}$$

In particular, curvatures $\mathrm{Rm}(g^i_{t_i})$ are uniformly bounded. Thus by the regularity of Ricci flow [Sh], $\|\mathrm{Rm}(g^i_{t_i})\|_{C^{k-2,\frac{1}{2}}(M,g_{t_i})}$ are uniformly bounded for any $k \ge 2$. Hence by Cheeger-Gromov's compactness theorem (cf. [GW]), one sees that there exist a subsequence $g^{i_k}_{t_{i_k}}$ of $g^i_{t_i}$ and a sequence of diffeomorphisms Φ_{i_k} on M such that $\Phi^\star_{i_k} g^{i_k}_{t_{i_k}}$ converge to a limit Riemannian metric g_∞ on M in the $C^{k,\frac{1}{2}}$-topology. Clearly by (5.4), it holds

$$2\epsilon_0 \ge \|g_\infty - g_{KE}\|_{C^3(M)} \ge \epsilon_0. \tag{5.5}$$

On the other hand, again by (5.4), the sequence of induced complex structures $(\Phi_{i_k}^{-1})_\star\cdot J\cdot(\Phi_{i_k})_\star$ (maybe after taking a subsequence of i_k) will converge to a $C^{k,\frac{1}{2}}$-differential complex structure J_∞ on M. By the regularity, J_∞ is analytic. Moreover, one can show that g_∞ is Kähler with respect to J_∞.

By the monotonicity of $\lambda(\cdot)$ along the flow (see (3.8) in Section 3), we have

$$\lambda(g^{i_k}) \le \lambda(g^{i_k}_{t_{i_k}}).$$

It follows

$$\lambda(g_\infty) \ge \lim_{i_k} \lambda(g^{i_k}) \ge \lambda(g_{KE}).$$

Thus by Lemma 3.4, we see that g_∞ is in fact a Kähler-Einstein metric on (M, J_∞). Now by Theorem 5.1, (g_∞, J_∞) is conjugate to (g_{KE}, J) by a diffeomorphism on M. But this contradicts to (5.5). The contradiction implies that (5.1) is true.

By (5.1), we have

$$\|g_t - g_{KE}\|_{C^3(M)} \le \epsilon_0 \le 1, \ \forall\ t > 0. \tag{5.6}$$

Thus, there exists a sequence $\{g_{t_i}\}$ of g_t which converges to a limit Riemannian metric g_∞ on M in the C^3-topology. Hence by Lemma 3.5, g_∞ is a Kähler-Ricci soliton. Moreover, by Corollary 4.6, $\lambda(g_\infty) = (2\pi)^{-n}nV = \lambda(g_{KE})$ since the K-energy is bounded from below [BM], [DT], and consequently, by Lemma 3.4, g_∞ must be a Kähler-Einstein metric. Therefore, by Theorem 5.1, we conclude that g_∞ is conjugate to g_{KE} by a diffeomorphism on M. Finally, The uniqueness of limit g_∞ enable us to prove that the KR-flow $(g_t; g)$ globally converges to g_{KE}. Proposition 2.1 is proved.

□

6. Proof of Theorem 2.3

Proof of Theorem 2.3. By Corollary 2.2, there exists a $\tau_0 \le 1$ with $[0, \tau_0) \subset I$. We need to show that $\tau_0 \in I$. In fact we want to prove that for any $\delta > 0$ there exists a large T such that

$$\|g^s_t - g_{KE}\|_{C^3(M)} \le \delta, \ \forall\ t \ge T \text{ and } s < \tau_0. \tag{6.1}$$

As we did for Proposition 2.1, we will prove it by contradiction. Suppose that we can find a sequence of Kähler metrics $g^{s_i}_{t_i}$, where $s_i \to \tau_0$ and $t_i \to \infty$, of KR-flows $(g^{s_i}_t; g^{s_i})$ and a sequence of diffeomorphisms Ψ_i on M such that

$$|\Psi_i^\star g^{s_i}_{t_i} - g_{KE}|_{C^3(M)} \ge \delta_0 > 0, \tag{6.2}$$

for some constant δ_0. Since the KR-flows $(g^{s_i}_t; g^{s_i})$ converge to g_{KE}, we may further assume that $g^{s_i}_{t_i}$ satisfy

$$|\Psi_i^\star g^{s_i}_{t_i} - g_{KE}|_{C^3(M)} \le 2\delta_0. \tag{6.3}$$

Then it follows from Cheeger-Gromov's compactness theorem that there is a subsequence $g_{t_{i_k}}^{s_{i_k}}$ of $g_{t_i}^i$ converging to a Kähler metric (g_∞, J_∞) with property

$$2\delta_0 \ge \|g_\infty - g_{KE}\|_{C^3(M)} \ge \delta_0. \tag{6.4}$$

We claim that (g_∞, J_∞) is a Kähler-Einstein metric. By Lemma 3.4, we only need to show that $\lambda(g_\infty) = \lambda(g_{KE}) = (2\pi)^{-n}nV$. It follows from Corollary 4.6 and the monotonicity of $\lambda(g_t^{\tau_0})$ that for any $\epsilon > 0$, there exists $T > 0$ such that

$$\lambda(g_t^{\tau_0}) \ge (2\pi)^{-n}nV - \frac{\epsilon}{2},\ \forall\ t \ge T.$$

Since Kähler-Ricci flow is stable in finite time and $\lambda(g_t^s)$ is monotonic in t, there is a small $\delta > 0$ such that for any $s \ge \tau_0 - \delta$, we have

$$\lambda(g_t^s) \ge (2\pi)^{-n}nV - \epsilon,\ \forall\ t \ge T. \tag{6.5}$$

Since $s_i \to \tau_0$ and $t_i \to \infty$, we conclude that

$$\lim_{s_i\to\tau_0, t_i\to\infty} \lambda(g_{t_i}^{s_i}) = (2\pi)^{-n}nV.$$

By the continuity of $\lambda(\cdot)$, it follows

$$\lambda(g_\infty) = (2\pi)^{-n}nV = \lambda(g_{KE}). \tag{6.6}$$

This proves the claim.

Now by Theorem 5.1, we see that (g_∞, J_∞) is conjugate to (g_{KE}, J) by a diffeomorphism on M. But this contradicts to (6.4). This contradiction implies that (6.1) is true.

By (6.1), we have

$$\|g_t^{\tau_0} - g_{KE}\|_{C^3(M)} \le \delta_0 \le 1,\ \forall\ t \ge T. \tag{6.7}$$

Thus, there exists a sequence $\{g_{t_i}^{\tau_0}\}$ of $g_t^{\tau_0}$ which converges to a limit Riemannian metric g_∞ on M in the C^3-topology. Hence, by Lemma 3.5, g_∞ is a Kähler-Ricci soliton. Moreover, by Corollary 4.6, $\lambda(g_\infty) = (2\pi)^{-n}nV$, so g_∞ must be a Kähler-Einstein metric. Therefore, by Theorem 5.1, we conclude that g_∞ is conjugate to g_{KE} by a diffeomorphism on M. Finally, the uniqueness of limit g_∞ enable us to prove that the KR-flow $(g_t^{\tau_0}; g^{\tau_0})$ globally converges to g_{KE}. This shows $\tau_0 \in I$.

□

Completion of proof of Theorem 1.1. We have shown that $I = [0, 1]$, so the first part of Theorem 1.1 is proved. The second part of Theorem 1.1 is known (cf. [PS], also see [CT2]). For the reader's convenience, we outline its proof. Since the limiting complex structure J_∞ is conjugate to J, by an estimate (3.11) in [PS], one sees that there exists two uniform constants $C, \delta >$ such that

$$\int_M |\nabla h_t|^2_{g_t}\omega^n_{g_t} \le Ce^{-\delta t},\ \forall\ t > 0. \tag{6.8}$$

Moreover, by using (6.8), one can further derive

$$\int_M |\nabla^p h_t|^2_{g_t}\omega^n_{g_t} \le C_p e^{-\delta_p t},\ \forall\ t > 0,$$

where $p > 0$ is any integer and two uniform constants $C_p, \delta_p > 0$ depend only on p (cf. [CT2], [PS], [Zhu], etc). By using the Sobolev embedding theorem, it follows

$$\|h_t - a\|_{C^0(M)} \le C' e^{-\delta' t},\ \forall\ t > 0,$$

for some constant a. As a consequence, we get

$$\|\phi_t - \frac{1}{V}\int_M \phi_t \omega^n_{g_{KE}}\|_{C^0(M)} \le C''.$$

Using this, we can prove that all C^ℓ-norms of ϕ_t decay exponentially. Hence, ϕ_t are convergent and the corresponding metrics ω_{ϕ_t} converge exponentially to a Kähler-Einstein metric g'_{KE} on (M, J). By the uniqueness theorem on Kähler-Einstein metrics, g'_{KE} must be conjugate to g_{KE} by a holomorphism transformation. The proof of Theorem 1.1 is completed.

□

7. Proof of Theorem 1.2

As in the proof of Theorem 1.1, we consider a path of Kähler forms $\omega_s = \omega_{g_0} + s\sqrt{-1}\partial\overline{\partial}\phi$ with $\omega_g = \omega_{g_0} + \sqrt{-1}\partial\overline{\partial}\phi$ and set

$$I = \{s \in\ [0,1]|\ \text{KR-flow}\ (g^s_t;\omega_s)\ \ \text{converges to}\ g_{KE}\ \ \text{in the}\ C^\infty\text{-topology}\}.$$

By the assumption, $0 \in I$. We shall show that I is both open and closed.

The following lemma should be due to Chen [Ch]. It generalizes Bando-Mabuchi's result about the lower bound of K-energy [BM].

Lemma 7.1. *If there exists a sequence of Kähler metrics on (M, J) with Kähler class $2\pi c_1(M)$ converging to a Kähler-Einstein metric g_{KE} for some complex structure J_0 in the C^∞-topology, then Mabuchi's K-energy is bounded from below on the space of Kähler potentials on (M, J).*

Proof. By our assumption, there are diffeomorphisms Ψ_i of M such that $|(\Psi_i^{-1})_\star \cdot J \cdot (\Psi_i)_\star - J_0|_{C^k(M)}$ converge to 0 ($\forall\ k > 0$). Then an observation of Szekylihidi in [Sz] implies that there exists a smooth holomorphic fibration: $\pi : Y \mapsto D$, where D denotes the unit disc in the complex plane, such that 1) $\pi^{-1}(z)$ is boholomorphic to (M, J); 2) $\pi^{-1}(0)$ is a compact Kähler manifold

(M, J') which admits a Kähler-Einstein metric g'_{KE}. In fact, by [CS], this J' is conjugate to J_0. Thus the K-energy is bounded from below on space of Kähler potentials on (M, J) by Chen's main result in [Ch]. □

The following proposition shows that I is an open set.

Proposition 7.2. *Let (M, J) be a compact Kähler manifold with $c_1(M) > 0$ and g_{KE} be a Kähler-Einstein metric with Kähler class $2\pi c_1(M)$ for some complex structure J' on M. Suppose that there exists a Kähler metric (J, g_0) with Kähler class $2\pi c_1(M)$ such that the KR-flow with the initial metric g_0 converges to g_{KE} in the C^3-topology. Then there exists $\delta > 0$ depending only on g_{KE} and g_0 such that for any Kähler metric g on (M, J) with Kähler class $2\pi c_1(M)$ satisfying*

$$\|g - g_0\|_{C^3(M)} \leq \delta, \tag{7.1}$$

the KR-flow $(g_t; g)$ converges to g_{KE} in the C^∞-topology.

Proof. Its proof is identical to that for Proposition 2.1. Since the KR-flow is stable in finite time, by our assumption, we see that for any $\epsilon > 0$, there exist $\delta > 0$ and a large T depending only on ϵ and g_0 such that for any Kähler metric g on (M, J) with Kähler class $2\pi c_1(M)$, whenever

$$\|g - g_0\|_{C^3(M)} \leq \delta,$$

then the evolved Kähler metric g_T of KR-flow $(g_t; g)$ satisfies

$$\|g_T - g_{KE}\|_{C^3(M)} < \epsilon. \tag{7.2}$$

We claim that (7.2) holds for any $t \geq T$. If the claim is false, there is a sequence of Kähler metrics g^i on (M, J) with Kähler class $2\pi c_1(M)$ satisfying

$$||g^i - g_0||_{C^3(M)} \to 0, \text{ as } i \to \infty, \tag{7.3}$$

such that there exist $t_i \to \infty$ satisfying:

$$2\epsilon \geq \|g^i_{t_i} - g_{KE}\|_{C^3(M)} \geq \epsilon, \tag{7.4}$$

where $g^i_{t_i}$ is the evolved metric of the KR-flow $(g^i_t; g^i)$ at time t_i. Then by Cheeger-Gromov's compactness theorem, there exists a subsequence $g^{i_k}_{t_{i_k}}$ of $g^i_{t_i}$ which converges to a Kähler metric (g_∞, J_∞) with property

$$2\epsilon \geq \|g_\infty - g_{KE}\|_{C^3(M)} \geq \epsilon. \tag{7.5}$$

Moreover, by the monotonicity of $\lambda(\cdot)$ and (7.3), it is easy to see

$$\lambda(g_\infty) \geq \lambda(g_{KE}).$$

It follows from Lemma 3.4 that g_∞ is actually a Kähler-Einstein metric on (M, J_∞). Consequently, by Theorem 5.1, (g_∞, J_∞) is conjugate to (g_{KE}, J')

by a diffeomorphism on M. But this contradicts to (7.5). So the claim is true.

By the claim, we see that

$$\|g_t - g_{KE}\|_{C^3(M)} \leq C, \tag{7.6}$$

for all Kähler metrics of $(g_t; g)$ as long as g satisfies (7.1). Then there exists a sequence $\{g_{t_i}\}$ of g_t which converges to a Kähler-Ricci soliton g_∞ according to Lemma 3.5. Moreover, since the K-energy is bounded from below according to Lemma 7.1, one can easily show that g_∞ is actually a Kähler-Einstein metric, which has to be conjugate to g_{KE} by a diffeomorphism of M. Therefore, the flow $(g_t; g)$ converges to g_{KE} in the C^∞-topology.

□

Proof of Theorem 1.2. By Proposition 7.2, we only need to prove that I is closed. However, the proof of Theorem 2.3 still applies here since Corollary 4.6 is still true by Lemma 7.1. So we refer the readers to Section 6 for showing that I is closed.

□

8. Appendix

In this appendix, we give a uniform L^∞-estimate for minimizing solutions of (3.5) associated to the evolved Kähler metrics of KR-flow. The result was used in Section 4 and is of independent interest.

Proposition 8.1. *Let $f = f_t$ be a minimizer of $W(g_t, \cdot)$ for the evolved Kähler metrics $g = g_t$ of (2.1). Then f is uniformly bounded.*

Proof. Let $u = e^{-\frac{f}{2}}$. Then by (3.5), u satisfies

$$\Delta u - \frac{1}{2} f u - \frac{1}{4} R u = \frac{1}{2}\lambda(g) u. \tag{8.1}$$

Since the scalar curvature R is uniformly bounded by Lemma 4.1, we have

$$-\Delta u \leq u^{1+\delta} + C_\delta, \tag{8.2}$$

for any $\delta > 0$. Thus for any $p > 1$, we have

$$-u^{\frac{p}{2}} \Delta u^{\frac{p}{2}} \leq \frac{p}{2} u^{p+\delta} + C_\delta \frac{p}{2} u^{p-1+\delta}.$$

By the integration by parts, we get

$$\begin{aligned}\int_M |\nabla u^{\frac{p}{2}}|^2 dV_g &\leq p\Big(\int_M u^{p+\delta} dV_g + C_1\Big)\\ &\leq p[(\|u\|_{L^{\frac{(1+\epsilon)\delta}{\epsilon}}})^{\frac{\epsilon}{1+\epsilon}})(\|u\|_{L^{(1+\epsilon)p}})^{\frac{1}{1+\epsilon}} + C_1]\\ &\leq C_2 p[(\|u\|_{L^{(1+\epsilon)p}})^{\frac{1}{1+\epsilon}} + 1], \end{aligned} \tag{8.3}$$

where $\epsilon < \frac{2}{n-2}$. At the last inequality we used the fact $\int_M u^2 = \int_M dV_g = V$. On the other hand, since the scalar curvature R is uniformly bounded, we have the Sobolev inequality (cf. [Zha]):

$$\int_M |\nabla\psi|^2 dV_g \geq \delta_0 \int_M \psi^{\frac{2n}{n-2}} dV_g - C_0 \int_M \psi^2 dV_g,$$

for any $W^{1,2}$-function ψ, where δ_0 and C_0 are uniform constants. In particular, for $\psi = u^{\frac{p}{2}}$, we have

$$\int_M |\nabla u^{\frac{p}{2}}|^2 dV_g \geq \delta_0 (\|u\|_{L^{\frac{np}{n-2}}})^{\frac{n-2}{n}} - C_0' \int_M u^p dV_g.$$

Thus we derive from (8.3)

$$\|u\|_{L^{\frac{np}{n-2}}} \leq (C_3 p)^{\frac{1}{p}} (\|u\|_{L^{(1+\epsilon)p}} + 1).$$

Iterating the above over p, we can deduce

$$\|u\|_{L^\infty} \leq C_4 (\int_M u^2 dV_g + 1) = C_4 (V+1),$$

consequently,

$$f \geq -C'. \tag{8.4}$$

Note the normalized condition $\int_M e^{-f} dV_g = V$. By (8.4), we can find an A and an open set E_A of M for each f such that

$$\int_{E_A} e^h dV_g > \delta,$$

where $E_A \subset \{x \in M | \, f(x) < A\}$ and $h = h_t$ is the Ricci potential function of $g = g_t$ with the normalization

$$\int_M e^h dV_g = V.$$

Note that h is uniformly bounded by Lemma 4.1. Hence, we have

$$\begin{aligned}\int_M f e^h dV_g &= \int_{M\setminus E_A} f e^h dV_g + \int_{E_A} f e^h dV_g \\ &\leq \left(\int_{M\setminus E_A} e^h dV_g\right)^{\frac{1}{2}} \left(\int_{M\setminus E_A} f^2 e^h dV_g\right)^{\frac{1}{2}} + CA.\end{aligned}$$

It follows

$$\int_M f^2 e^h dV_g \geq \frac{1}{V-\delta} (\int_M f e^h dV_g)^2 - a(\int_M f^2 e^h dV_g)^{\frac{1}{2}} - C', \tag{8.5}$$

for some constant $a, C' > 0$. On the other hand, by integrating (3.5) at both two sides, we have

$$\int_M |\nabla f|^2 dV_g \leq \int_M f dV_g + C''.$$

Then by using the following weighted Poincare's inequality (cf. [TZ3]),

$$\int_M |\nabla f|^2 e^h dV_g \geq \int_M f^2 e^h dV_g - \frac{1}{V}(\int_M f e^h dV_g)^2,$$

we obtain

$$\int_M f^2 e^h dV_g \leq \frac{1}{V}(\int_M f e^h dV_g)^2 + b\int_M f dV_g + C''', \tag{8.6}$$

for some constant $b, C''' > 0$. Combining this with (8.5), we deduce

$$\int_M f^2 e^h dV_g \leq C_6,$$

consequently,

$$\int_M f^2 dV_g \leq C_7. \tag{8.7}$$

It follows from (3.5) that f satisfies,

$$\Delta f \geq -f - B.$$

Then by the standard L^∞-estimate for elliptic equations of second order, we have

$$\|f^+\|_{L^\infty} \leq C(\int_M f^2 dV_g + 1) \leq C_8,$$

where $f^+ = \max\{f, 0\}$. Thus we have proved that f is uniformly bounded. □

References

[1] [Ba] Bando, S., On three dimemsional compact Kähler manifolds of nonnegative bisectional curvature, J. Differ. Geom., 19 (1984), 284-297.

[2] [BM] Bando, S. and Mabuchi, T., Uniqueness of Kähler Einstein metrics modulo connected group actions, Algebraic Geometry, Adv. Stud. Pure Math. (1987), 11-40.

[3] [Ca] Cao, H.D., Deformation of Kähler metrics to Kähler-Einstein metrics on compact Kähler manifolds, Invent. Math., 81 (1985), 359-372.

[4] [CS] Chen, X.X. and Sun, S., Calabi flow, geodesic rays, and uniqueness of constant scalar curvature Kähler metrics, arXiv:1004.2012v1, 2010.

[5] [Ch] Chen, X.X., The space of Kähler metrics (IV)-On the lower bound of K-energy, arXiv:0809.4081, 2008.

[6] [CT1] Chen, X.X. and Tian, G., Ricci flow on Kähler-Einstein surfaces, Invent. Math., 147 (2002), 487-544.

[7] [CT2] Chen, X.X. and Tian, G., Ricci flow on Kähler-Einstein manifolds, Duke Math. J., 131 (2006), 17-73.

[8] [DT] Ding, W. and Tian, G., The generalized Moser-Trudinger inequality, in Nonlinear Analysis and Microlocal Analysis: Proceedings of the International Conference at Nakai Institue of Mathematics (K-C, Chang et., Eds), World Scientific, 1992, 57-70.

[9] [GW] Greene, R., Wu, H., Lipschitz convergence of Riemannian manifolds, Pacific J. Math., 131 (1988), 119-141.

[10] [Ma] Mabuchi, T., Some symplectic geometry on compact Kähler manifolds, I, Osaka J. Math., 24 (1987), 227-252.

[11] [Pe] Perelman, G., The entropy formula for the Ricci flow and its geometric applications, Arxiv/0211159.

[12] [PS] Phong, D.H., Song, J., On stability and convergence of Kähler-Ricci flow, J. Differ. Geom., 72 (2006), no 1, 149-168.

[13] [PSSW] Phong, D.H., Song, J., Sturm, J. and Weinkove, B., The Kähler-Ricci flow and $\overline{\partial}$-operator on vector fields, J. Differ. Geom., 81 (2009), 631-647.

[14] [Ro] Rothaus, O., Logarithmic Sobolev inequality and the spectrum of Schrödinger operators, J. Funct. Analysis, 42 (1981), no. 1, 110-120.

[15] [Sh] Shi, W. X., Ricci deformation of the metric on complete noncopmact Riemannian manifolds, J. Diff. Geom., 30 (1989), 223-301.

[16] [ST] Sesum, N. and Tian, G., Bounding scalar curvature and diameter along the Kähler-Ricci flow (after Perelman), J. Inst. Math., Jussieu, 7 (2008), 575-587.

[17] [SW] Sun, S. and Wang, Y., On Kähler-Ricci flow near a Kähler-Einstein metric, arXiv:1004.2018, 2010.

[18] [Sz] Szekelyhidi, G., The Kähler-Ricci flow and K-polystability, Amer. J. Math., 132 (2010), 1077-1090.

[19] [T1] Tian, G., Kähler-Einstein metrics with positive scalar curvature, Invent. Math., 101 (1990), 101-172.

[20] [T2] Tian, G., Extremal metrics and geometric stability, Special issue for S. S. Chern, Houston J. Math., 28 (2002), no 2, 411-432.

[21] [TZ1] Tian, G. and Zhu, X.H., A nonliner inequality of Moser-Trudinger type, Calculus of Variations, 10 (2000), 349-354.

[22] [TZ2] Tian, G. and Zhu, X., Uniqueness of Kähler-Ricci solitons, Acta Math., 184 (2000), 271-305.

[23] [TZ3] Tian, G. and Zhu, X., Convergence of the Kähler-Ricci flow, J. Amer Math. Sci., 17 (2006), 675-699.

[24] [TZ4] Tian, G. and Zhu, X., Perelman's W-functional and stability of Kähler-Ricci flow, Arxiv/0801.3504.

[25] [Zha] Zhang, Q. S., A uniform Sobolev inequality under Ricci flow, Arxiv:0706.1329.

[26] [Zhu] Zhu, X., Stability on Kähler-Ricci flow on a compact Kähler manifold with positive first Chern class; I, Arxiv/0908.1488v1.

Gang Tian, School of Mathematical Sciences and BICMR, Peking University, Beijing, 100871, China, and Department of Mathematics, Princeton University, New Jersey, NJ 02139, USA, tian@math.mit.edu

Xiaohua Zhu, School of Mathematical Sciences and BICMR, Peking University, Beijing, 100871, China, xhzhu@math.pku.edu.cn